



1740

Solutio problematum quorundam astronomicorum

Leonhard Euler

Follow this and additional works at: <https://scholarlycommons.pacific.edu/euler-works>

 Part of the [Mathematics Commons](#)

Record Created:

2018-09-25

Recommended Citation

Euler, Leonhard, "Solutio problematum quorundam astronomicorum" (1740). *Euler Archive - All Works*. 39.
<https://scholarlycommons.pacific.edu/euler-works/39>

This Article is brought to you for free and open access by the Euler Archive at Scholarly Commons. It has been accepted for inclusion in Euler Archive - All Works by an authorized administrator of Scholarly Commons. For more information, please contact mgibney@pacific.edu.

SOLVTIO PROBLEMATVM QVORVNDAM ASTRONOMICORVM.

AVCTORE

Leonb. Eulero.

Problema 1.

Data planetae aequatione maxima, inuenire orbitae eius excentricitatem.

Solutio.

Conuertatur aequatio maxima in minuta secunda, sitque eorum numerus = m ; dico fore distantiam planetae a Sole mediam ad excentricitatem vt 412533 ad numerum m ; si quidem aequatio non fuerit nimis magna. At si aequatio admodum fuerit ingens, posita ratione distantiae mediae ad excentricitatem vt 1 ad v , erit $v = \frac{m}{412533} - \frac{m^2}{32(412533)^2}$. Q.E.I.

Problema 2:

Data excentricitate orbitae planetaris, inuenire aequationem maximam.

Solutio.

Sit 1 ad v vt distantia planetae a Sole media ad excentricitatem, et sit m numerus minorum secundorum aequationis maximae, qui quaeritur, dico fore $m = 412533 (v + \frac{v^2}{32})$; vel per logarithmos erit $\log. m =$
Tom. VII. N 5,

98 SOLVTIO PROBLEMATVM QVORVND. ASTR.

$5,6154596 + l(v + \frac{v^7}{32})$. Vbi notandum est, nisi excentricitas fuerit vehementer magna loco quantitatis $v + \frac{v^7}{32}$ sumi posse tantum v . Q. E. I.

Problema 3.

Data excentricitate orbitae planetaris, inuenire anomaliam mediam, cui aequatio maxima respondet.

Solutio.

Sit r ad v vt distantia media ad excentricitatem, quae ergo ratio datur et proinde v . Multiplicetur sinus totus per $\frac{v^3}{4}$, et factum erit sinus cuiusdam anguli ex tabulis inueniendi: fit hic angulus q graduum. *Haec vero operatio commodius per logarithmos instituetur.*

Deinde quaeratur logarithmus quantitatis $v - \frac{v^7}{32}$, vel tantum ipsius v , ex tabulis logarith. num. naturalium, si fuerit v admodum paruum, iste logarithmus addatur ad hunc $5,3144295$, et logarithmi, qui prodit, quaeratur numerus respondens, qui fit n ; vbi notetur n'' esse dimidiam partem aequationis maximae; ita vt, si aequatio maxima iam fuerit inuenta hac posteriori operatione nequidem sit opus. Dico fore anomaliam mediam quaesitam $90^\circ + q^\circ + n''$. Q. E. I.