



Spring 4-17-2015

Euler's Multiple Solutions to a Diophantine Problem

Christopher D. Goff

University of the Pacific, cgoфф@pacific.edu

Follow this and additional works at: <https://scholarlycommons.pacific.edu/cop-facpres>

 Part of the [Mathematics Commons](#)

Recommended Citation

Goff, C. D. (2015). Euler's Multiple Solutions to a Diophantine Problem. Paper presented at CMC³ Recreational Math Conference in Lake Tahoe, CA.

<https://scholarlycommons.pacific.edu/cop-facpres/474>

This Lecture is brought to you for free and open access by the All Faculty Scholarship at Scholarly Commons. It has been accepted for inclusion in College of the Pacific Faculty Presentations by an authorized administrator of Scholarly Commons. For more information, please contact mgibney@pacific.edu.

Euler's Multiple Solutions to a Diophantine Problem

Christopher Goff

University of the Pacific

18 April 2015

Leonhard Euler (1707-1783)



- Swiss
- Had 13 kids
- Worked in St. Petersburg and Berlin
- By 1735, blind in right eye – went totally blind later, but kept writing (secretary)
- Published 530 books and papers in his life, and many more after his death (including the ones we will consider)
- Very prolific and successful, but also not always rigorous

Graphic from <http://sebastianiaguirre.wordpress.com/2011/04/12/project-euler/>

Some of Euler's Mathematics

- 1 Using certain notations: $f(x)$, e , \sum , i
- 2 Using a, b, c for the sides of a right triangle
- 3 $e^{ix} = \cos x + i \sin x$ [$e^{i\pi} + 1 = 0$]
- 4 $V - E + F = 2$, Ex: cube (8 vertices, 12 edges, 6 faces)

More of Euler's Mathematics

$$① \quad \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1} + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots = \frac{\pi^2}{6}$$

- ② Euler line (geometry)
- ③ Euler's method (ordinary differential equations)
- ④ Eulerian path (graph theory)

A Problem


- Find two positive integers satisfying the following properties.
 - Their sum is a square of an integer.
 - The sum of their squares is the fourth power of an integer.

A Problem

- Find two positive integers satisfying the following properties.
 - Their sum is a square of an integer.
 - The sum of their squares is the fourth power of an integer.
- “On finding three or more numbers, the sum of which is a square and the sum of the squares of which is a fourth power” (1824).

A Problem

- Find two positive integers satisfying the following properties.
 - Their sum is a square of an integer.
 - The sum of their squares is the fourth power of an integer.
- “On finding three or more numbers, the sum of which is a square and the sum of the squares of which is a fourth power” (1824).
- Objective: understand Euler’s solution and follow his algebraic twists and turns along the way.



Where did this problem come from?

- Diophantus (c.200 - c.284) of Alexandria and “Diophantine”

Where did this problem come from?

- Diophantus (c.200 - c.284) of Alexandria and “Diophantine”
- *Find a right triangle whose legs add up to a perfect square and whose hypotenuse is also a perfect square.*

Where did this problem come from?

- Diophantus (c.200 - c.284) of Alexandria and “Diophantine”
- *Find a right triangle whose legs add up to a perfect square and whose hypotenuse is also a perfect square.* That is, if the legs are x and y , and the hypotenuse is $z = N^2$, then

$$x + y = M^2 \quad \text{and} \quad x^2 + y^2 = z^2 = N^4.$$

Where did this problem come from?

- Diophantus (c.200 - c.284) of Alexandria and “Diophantine”
- *Find a right triangle whose legs add up to a perfect square and whose hypotenuse is also a perfect square.* That is, if the legs are x and y , and the hypotenuse is $z = N^2$, then

$$x + y = M^2 \quad \text{and} \quad x^2 + y^2 = z^2 = N^4.$$

- Also solved by Pierre de Fermat (1601-1665) and Joseph-Louis Lagrange (1736-1813), before Euler.

Where did this problem come from?

- Diophantus (c.200 - c.284) of Alexandria and “Diophantine”
- *Find a right triangle whose legs add up to a perfect square and whose hypotenuse is also a perfect square.* That is, if the legs are x and y , and the hypotenuse is $z = N^2$, then

$$x + y = M^2 \quad \text{and} \quad x^2 + y^2 = z^2 = N^4.$$

- Also solved by Pierre de Fermat (1601-1665) and Joseph-Louis Lagrange (1736-1813), before Euler.
- Which is bigger, M or N ? Why?

My translations are NOT literal, but get the point across.

Patterns and facts

- $(s + t)^2 = s^2 + 2st + t^2$
- $(s + t + u)^2 = s^2 + t^2 + u^2 + 2st + 2su + 2tu$
- In the quadratic $ax^2 + bx + c = 0$, the sum of the two roots is $-\frac{b}{a}$.

Pythagorean Triples (a, b, c)

- Euclid (c.300 BCE): $a = p^2 - q^2$; $b = 2pq$; $c = p^2 + q^2$.

Pythagorean Triples (a, b, c)

- Euclid (c.300 BCE): $a = p^2 - q^2$; $b = 2pq$; $c = p^2 + q^2$.

$$a^2 + b^2 = (p^2 - q^2)^2 + (2pq)^2$$

Pythagorean Triples (a, b, c)

- Euclid (c.300 BCE): $a = p^2 - q^2$; $b = 2pq$; $c = p^2 + q^2$.

$$\begin{aligned}a^2 + b^2 &= (p^2 - q^2)^2 + (2pq)^2 \\ &= p^4 - 2p^2q^2 + q^4 + 4p^2q^2\end{aligned}$$

Pythagorean Triples (a, b, c)

- Euclid (c.300 BCE): $a = p^2 - q^2$; $b = 2pq$; $c = p^2 + q^2$.

$$\begin{aligned}a^2 + b^2 &= (p^2 - q^2)^2 + (2pq)^2 \\ &= p^4 - 2p^2q^2 + q^4 + 4p^2q^2 \\ &= p^4 + 2p^2q^2 + q^4\end{aligned}$$

Pythagorean Triples (a, b, c)

- Euclid (c.300 BCE): $a = p^2 - q^2$; $b = 2pq$; $c = p^2 + q^2$.

$$\begin{aligned}a^2 + b^2 &= (p^2 - q^2)^2 + (2pq)^2 \\&= p^4 - 2p^2q^2 + q^4 + 4p^2q^2 \\&= p^4 + 2p^2q^2 + q^4 \\&= (p^2 + q^2)^2 = c^2. \quad \square\end{aligned}$$

Pythagorean Triples (a, b, c)

- Euclid (c.300 BCE): $a = p^2 - q^2$; $b = 2pq$; $c = p^2 + q^2$.

$$\begin{aligned}a^2 + b^2 &= (p^2 - q^2)^2 + (2pq)^2 \\&= p^4 - 2p^2q^2 + q^4 + 4p^2q^2 \\&= p^4 + 2p^2q^2 + q^4 \\&= (p^2 + q^2)^2 = c^2. \quad \square\end{aligned}$$

- Euclid showed that EVERY primitive Pythagorean triple can be put into this form, for some choice of p and q .

Euler's solution to Diophantus' problem: §5

Find x, y so that $x + y = M^2$ and $x^2 + y^2 = N^4$.

Euler's solution to Diophantus' problem: §5

Find x, y so that $x + y = M^2$ and $x^2 + y^2 = N^4$.

- “Let us begin with the second condition. First, the formula $xx + yy$ shall be made a square, by placing $x = aa - bb$ and $y = 2ab$, for then $xx + yy = (aa + bb)^2$. [**]

Euler's solution to Diophantus' problem: §5

Find x, y so that $x + y = M^2$ and $x^2 + y^2 = N^4$.

- “Let us begin with the second condition. First, the formula $xx + yy$ shall be made a square, by placing $x = aa - bb$ and $y = 2ab$, for then $xx + yy = (aa + bb)^2$. [**]
- In addition, $a^2 + b^2$ should be a square, which happens in the same way by setting $a = p^2 - q^2$ and $b = 2pq$: from here, it follows that $x^2 + y^2 = (a^2 + b^2)^2 = (p^2 + q^2)^4$, and thus the latter condition has now been fully satisfied. [**]

Euler's solution to Diophantus' problem: §5

Find x, y so that $x + y = M^2$ and $x^2 + y^2 = N^4$.

- “Let us begin with the second condition. First, the formula $xx + yy$ shall be made a square, by placing $x = aa - bb$ and $y = 2ab$, for then $xx + yy = (aa + bb)^2$. [**]
- In addition, $a^2 + b^2$ should be a square, which happens in the same way by setting $a = p^2 - q^2$ and $b = 2pq$: from here, it follows that $x^2 + y^2 = (a^2 + b^2)^2 = (p^2 + q^2)^4$, and thus the latter condition has now been fully satisfied. [**]
- Then, it remains to satisfy the first condition, namely that $x + y$ be a square.”

Euler's solution: §6

- “From these facts it is found that

$$x = a^2 - b^2 = p^4 - 6p^2q^2 + q^4 \quad \text{and} \quad y = 2ab = 4p^3q - 4pq^3;$$

and so the following formula $[x + y]$ ought to be a square

$$p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4, \dots$$

[with $p > q > 0$ and $a > b$].”

Euler's solution: §6

- “From these facts it is found that

$$x = a^2 - b^2 = p^4 - 6p^2q^2 + q^4 \quad \text{and} \quad y = 2ab = 4p^3q - 4pq^3;$$

and so the following formula $[x + y]$ ought to be a square

$$p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4, \dots$$

[with $p > q > 0$ and $a > b$].”

- Why do we have to pick $p > q$? Why do we have to have $a > b$?

Euler's solution: §7

- “The formula is solved by setting $\sqrt{x+y} = p^2 - 2pq + q^2$, from which $\frac{p}{q} = \frac{3}{2}$, or $p = 3$ and $q = 2$.”

Euler's solution: §7

- “The formula is solved by setting $\sqrt{x+y} = p^2 - 2pq + q^2$, from which $\frac{p}{q} = \frac{3}{2}$, or $p = 3$ and $q = 2$.”
- Wait a minute. What is Euler doing? How did he get that?

Euler's solution: §7

- “The formula is solved by setting $\sqrt{x + y} = p^2 - 2pq + q^2$, from which $\frac{p}{q} = \frac{3}{2}$, or $p = 3$ and $q = 2$.”
- Wait a minute. What is Euler doing? How did he get that?
- His answer is close to correct, but let's see how close. Let's square it.

Euler's solution: §7

- “The formula is solved by setting $\sqrt{x+y} = p^2 - 2pq + q^2$, from which $\frac{p}{q} = \frac{3}{2}$, or $p = 3$ and $q = 2$.”
- Wait a minute. What is Euler doing? How did he get that?
- His answer is close to correct, but let's see how close. Let's square it.

$$(p^2 - 2pq + q^2)^2 = p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4,$$

which doesn't quite equal $p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4$, as he claimed.

Euler's solution: §7 (cont.)

But he is close. Three of the terms are identical, and the other two just have different signs. So, let's set the two expressions equal and see what happens.

Euler's solution: §7 (cont.)

But he is close. Three of the terms are identical, and the other two just have different signs. So, let's set the two expressions equal and see what happens.

$$p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4 = p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4$$

Euler's solution: §7 (cont.)

But he is close. Three of the terms are identical, and the other two just have different signs. So, let's set the two expressions equal and see what happens.

$$\begin{aligned} p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4 &= p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4 \\ 12p^2q^2 &= 8p^3q \end{aligned}$$

Euler's solution: §7 (cont.)

But he is close. Three of the terms are identical, and the other two just have different signs. So, let's set the two expressions equal and see what happens.

$$\begin{aligned}p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4 &= p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4 \\12p^2q^2 &= 8p^3q \\3q &= 2p, \text{ or } \frac{p}{q} = \frac{3}{2}.\end{aligned}$$

Euler doesn't need the formula to be a square ALWAYS (like an identity). He ONLY needs to find values of p and q that make the formula equal to a square.

Euler's solution: §7 (cont.)

But he is close. Three of the terms are identical, and the other two just have different signs. So, let's set the two expressions equal and see what happens.

$$\begin{aligned}p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4 &= p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4 \\12p^2q^2 &= 8p^3q \\3q &= 2p, \text{ or } \frac{p}{q} = \frac{3}{2}.\end{aligned}$$

Euler doesn't need the formula to be a square ALWAYS (like an identity). He ONLY needs to find values of p and q that make the formula equal to a square.

- “The formula is solved by setting $\sqrt{x+y} = p^2 - 2pq + q^2$, from which $\frac{p}{q} = \frac{3}{2}$, or $p = 3$ and $q = 2$. [**]”

Euler's solution: §7 (cont.)

But he is close. Three of the terms are identical, and the other two just have different signs. So, let's set the two expressions equal and see what happens.

$$\begin{aligned}p^4 - 4p^3q + 6p^2q^2 - 4pq^3 + q^4 &= p^4 + 4p^3q - 6p^2q^2 - 4pq^3 + q^4 \\12p^2q^2 &= 8p^3q \\3q &= 2p, \text{ or } \frac{p}{q} = \frac{3}{2}.\end{aligned}$$

Euler doesn't need the formula to be a square ALWAYS (like an identity). He ONLY needs to find values of p and q that make the formula equal to a square.

- “The formula is solved by setting $\sqrt{x+y} = p^2 - 2pq + q^2$, from which $\frac{p}{q} = \frac{3}{2}$, or $p = 3$ and $q = 2$. [**]
- But then $a = 5$, and $b = 12$, and so $x < 0$, and this solution is rejected.” [$x = -119$; $y = 120$]

More of Euler's Algebra Skills

“On account of this, a new method must be established . . . and so we keep $q = 2$ but at the same time we put $p = 3 + v$, from which we deduce the following values:

More of Euler's Algebra Skills

“On account of this, a new method must be established . . . and so we keep $q = 2$ but at the same time we put $p = 3 + v$, from which we deduce the following values:

$$\begin{aligned}p^4 &= 81 + 108v + 54v^2 + 12v^3 + v^4, \\4p^3q &= 216 + 216v + 72v^2 + 8v^3, \\6p^2q^2 &= 216 + 144v + 24v^2, \\4pq^3 &= 96 + 32v, \\q^4 &= 16.\end{aligned}$$

When the terms are collected, the . . . formula adopts this form:”

More of Euler's Algebra Skills

“On account of this, a new method must be established . . . and so we keep $q = 2$ but at the same time we put $p = 3 + v$, from which we deduce the following values:

$$\begin{aligned}p^4 &= 81 + 108v + 54v^2 + 12v^3 + v^4, \\4p^3q &= 216 + 216v + 72v^2 + 8v^3, \\6p^2q^2 &= 216 + 144v + 24v^2, \\4pq^3 &= 96 + 32v, \\q^4 &= 16.\end{aligned}$$

When the terms are collected, the . . . formula adopts this form:”

$$x + y = 1 + 148v + 102v^2 + 20v^3 + v^4$$

More of Euler's Algebra Skills

“On account of this, a new method must be established . . . and so we keep $q = 2$ but at the same time we put $p = 3 + v$, from which we deduce the following values:

$$\begin{aligned}p^4 &= 81 + 108v + 54v^2 + 12v^3 + v^4, \\4p^3q &= 216 + 216v + 72v^2 + 8v^3, \\6p^2q^2 &= 216 + 144v + 24v^2, \\4pq^3 &= 96 + 32v, \\q^4 &= 16.\end{aligned}$$

When the terms are collected, the . . . formula adopts this form:”

$$x + y = 1 + 148v + 102v^2 + 20v^3 + v^4$$

Euler then guesses the square root of this to be: $1 + 74v - v^2$. [Why?](#)

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

$$(1 + 74v - v^2)^2 = 1 + 148v + 5474v^2 - 148v^3 + v^4,$$

which is not quite $1 + 148v + 102v^2 + 20v^3 + v^4$, but three of the terms are identical.

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

$$(1 + 74v - v^2)^2 = 1 + 148v + 5474v^2 - 148v^3 + v^4,$$

which is not quite $1 + 148v + 102v^2 + 20v^3 + v^4$, but three of the terms are identical. So, when setting them equal, several terms cancel, leaving:

$$102v^2 + 20v^3 = 5474v^2 - 148v^3$$

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

$$(1 + 74v - v^2)^2 = 1 + 148v + 5474v^2 - 148v^3 + v^4,$$

which is not quite $1 + 148v + 102v^2 + 20v^3 + v^4$, but three of the terms are identical. So, when setting them equal, several terms cancel, leaving:

$$\begin{aligned}102v^2 + 20v^3 &= 5474v^2 - 148v^3 \\168v^3 &= 5372v^2\end{aligned}$$

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

$$(1 + 74v - v^2)^2 = 1 + 148v + 5474v^2 - 148v^3 + v^4,$$

which is not quite $1 + 148v + 102v^2 + 20v^3 + v^4$, but three of the terms are identical. So, when setting them equal, several terms cancel, leaving:

$$\begin{aligned}102v^2 + 20v^3 &= 5474v^2 - 148v^3 \\168v^3 &= 5372v^2 \\v &= \frac{5372}{168} = \frac{1343}{42}.\end{aligned}$$

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

$$(1 + 74v - v^2)^2 = 1 + 148v + 5474v^2 - 148v^3 + v^4,$$

which is not quite $1 + 148v + 102v^2 + 20v^3 + v^4$, but three of the terms are identical. So, when setting them equal, several terms cancel, leaving:

$$\begin{aligned}102v^2 + 20v^3 &= 5474v^2 - 148v^3 \\168v^3 &= 5372v^2 \\v &= \frac{5372}{168} = \frac{1343}{42}.\end{aligned}$$

So $p = 3 + v = \frac{1469}{42}$ and $q = 2$. But Euler knew that he could multiply p and q by any constant and still have a perfect square.

More of Euler's Algebra Skills (cont.)

Answer: it's the same idea as before.

$$(1 + 74v - v^2)^2 = 1 + 148v + 5474v^2 - 148v^3 + v^4,$$

which is not quite $1 + 148v + 102v^2 + 20v^3 + v^4$, but three of the terms are identical. So, when setting them equal, several terms cancel, leaving:

$$\begin{aligned}102v^2 + 20v^3 &= 5474v^2 - 148v^3 \\168v^3 &= 5372v^2 \\v &= \frac{5372}{168} = \frac{1343}{42}.\end{aligned}$$

So $p = 3 + v = \frac{1469}{42}$ and $q = 2$. But Euler knew that he could multiply p and q by any constant and still have a perfect square. So he gets

$$p = 1469 \quad \text{and} \quad q = 84.$$

The Answer to Diophantus' Problem

Working backwards, Euler now gets

The Answer to Diophantus' Problem

Working backwards, Euler now gets

- $a = p^2 - q^2 = 1469^2 - 84^2 = 2,150,905$

The Answer to Diophantus' Problem

Working backwards, Euler now gets

- $a = p^2 - q^2 = 1469^2 - 84^2 = 2,150,905$
- $b = 2pq = 2(1469)(84) = 246,792$

The Answer to Diophantus' Problem

Working backwards, Euler now gets

- $a = p^2 - q^2 = 1469^2 - 84^2 = 2,150,905$
- $b = 2pq = 2(1469)(84) = 246,792$
- $x = a^2 - b^2 = 4,565,486,027,761$ and

The Answer to Diophantus' Problem

Working backwards, Euler now gets

- $a = p^2 - q^2 = 1469^2 - 84^2 = 2,150,905$
- $b = 2pq = 2(1469)(84) = 246,792$
- $x = a^2 - b^2 = 4,565,486,027,761$ and
- $y = 2ab = 1,061,652,293,520,$

The Answer to Diophantus' Problem

Working backwards, Euler now gets

- $a = p^2 - q^2 = 1469^2 - 84^2 = 2,150,905$
- $b = 2pq = 2(1469)(84) = 246,792$
- $x = a^2 - b^2 = 4,565,486,027,761$ and
- $y = 2ab = 1,061,652,293,520,$

“... which are the same that Fermat, and others after him, found. The sum of them is the square of the number 2,372,159, while the sum of the squares is the fourth power of the number 2,165,017.”

The Answer to Diophantus' Problem

Working backwards, Euler now gets

- $a = p^2 - q^2 = 1469^2 - 84^2 = 2,150,905$
- $b = 2pq = 2(1469)(84) = 246,792$
- $x = a^2 - b^2 = 4,565,486,027,761$ and
- $y = 2ab = 1,061,652,293,520,$

“... which are the same that Fermat, and others after him, found. The sum of them is the square of the number 2,372,159, while the sum of the squares is the fourth power of the number 2,165,017.”

WOW!!! But wait, there's more

Finding a Pattern, I

Next, Euler finds three numbers (x, y, z) .

- 1 Set $x = a^2 + b^2 - c^2$, $y = 2ac$, $z = 2bc$.
- 2 Then set $a = p^2 + q^2 - r^2$, $b = 2pr$, $c = 2qr$.
- 3 Then GUESS the square root of $x + y + z$.
- 4 ... Euler finds $p = r + \frac{3}{2}q$.
- 5 He then chooses $q = 2$, $r = 1$ to get $p = 4$ and thus ...
- 6 "x = 409; y = 152; z = 64, the sum of which is $x + y + z = 625 = 25^2$; while the sum of the squares will be $xx + yy + zz = 194,481 = 441^2 = 21^4$." (MUCH SMALLER)

Finding a Pattern, II

Next, Euler finds four numbers (x, y, z, v) .

- 1 Set $x = a^2 + b^2 + c^2 - d^2$, $y = 2ad$, $z = 2bd$, $v = 2cd$.
- 2 Then set $a = p^2 + q^2 + r^2 - s^2$, $b = 2ps$, $c = 2qs$, $d = 2rs$.
- 3 Then GUESS the square root of $x + y + z + v$.
- 4 ... Euler finds $p = s + \frac{3}{2}r - q$.
- 5 He then chooses $r = 2$, $q = s = 1$ to get $p = 3$ and thus ...
- 6 "x = 193; y = 104; z = 48; v = 16, the sum of which is $x + y + z + v = 361 = 19^2$; while the sum of the squares will be $xx + yy + zz + vv = (pp + qq + rr + ss)^4 = 15^4$."

Finding a Pattern, III

Next, Euler finds five numbers (x, y, z, v, w) .

- 1 Set $x = a^2 + b^2 + c^2 + d^2 - e^2$, $y = 2ae$, $z = 2be$, $v = 2ce$,
 $w = 2de$.
- 2 Then set $a = p^2 + q^2 + r^2 + s^2 - t^2$, $b = 2pt$, $c = 2qt$, $d = 2rt$,
 $e = 2st$.
- 3 Then GUESS the square root of $x + y + z + v + w$.
- 4 ... Euler finds $p = t + \frac{3}{2}s - r - q$.
- 5 He then chooses $s = 2$, $t = r = q = 1$ to get $p = 2$ and thus ...
- 6 "x = 89; y = 72; z = 32; v = 16; w = 16, the sum of which is
 $x + y + z + v + w = 225 = 15^2$; while the sum of the squares will be
 $x^2 + y^2 + z^2 + v^2 + w^2 = 11^4$."

The Pattern

The Pattern

- For 3 numbers, $p = r + \frac{3}{2}q$.

The Pattern

- For 3 numbers, $p = r + \frac{3}{2}q$.
- For 4 numbers, $p = s + \frac{3}{2}r - q$.

The Pattern

- For 3 numbers, $p = r + \frac{3}{2}q$.
- For 4 numbers, $p = s + \frac{3}{2}r - q$.
- For 5 numbers, $p = t + \frac{3}{2}s - r - q$.

The Pattern

- For 3 numbers, $p = r + \frac{3}{2}q$.
- For 4 numbers, $p = s + \frac{3}{2}r - q$.
- For 5 numbers, $p = t + \frac{3}{2}s - r - q$.
You try it!!
- For 6 numbers,

The Pattern

- For 3 numbers, $p = r + \frac{3}{2}q$.
- For 4 numbers, $p = s + \frac{3}{2}r - q$.
- For 5 numbers, $p = t + \frac{3}{2}s - r - q$.
You try it!!
- For 6 numbers, $p = u + \frac{3}{2}t - s - r - q$.

Student Work

- I teach Topics in the History of Mathematics. I assign a project in which students have to engage with a primary source or a translation of a primary source.
- One student chose this paper.
- She found six numbers that had the same property. Namely: 97, 112, 64, 64, 64, and 128.
- Their sum is $529 = 23^2$, and the sum of their squares is $50625 = 15^4$.

Another solution

“On a notable advancement in Diophantine analysis” (1830) has a different solution. [Why?](#)

Another solution

“On a notable advancement in Diophantine analysis” (1830) has a different solution. [Why?](#)

... because Lagrange criticized Euler's original solution method. So Euler wrote two more papers going into more generality about how to generate solutions.

Euler generalizes to find integer solutions to

$$a^2x^4 + 2abx^3y + cx^2y^2 + 2bdxy^3 + d^2y^4 = \square$$

by making substitutions and taking advantage of certain patterns. We'll work through an example.

Euler generalizes to find integer solutions to

$$a^2x^4 + 2abx^3y + cx^2y^2 + 2bdxy^3 + d^2y^4 = \square$$

by making substitutions and taking advantage of certain patterns. We'll work through an example.

But first, we'll set $y = 1$ and look for rational solutions. [Why is this OK?](#)

A Related Example: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$

A Related Example: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$

This is rewritten as: $(1 + 6x + x^2)^2 - 32x^2 = \square$. If we let

$$1 + 6x + x^2 = p^2 + 8q^2 \quad \text{and} \quad x = pq,$$

then

$$(1 + 6x + x^2)^2 - 32x^2 = (p^2 - 8q^2)^2.$$

A Related Example: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$

This is rewritten as: $(1 + 6x + x^2)^2 - 32x^2 = \square$. If we let

$$1 + 6x + x^2 = p^2 + 8q^2 \quad \text{and} \quad x = pq,$$

then

$$(1 + 6x + x^2)^2 - 32x^2 = (p^2 - 8q^2)^2.$$

So now we have to find solutions to:

$$1 + 6pq + p^2q^2 = p^2 + 8q^2,$$

which is quadratic in p or in q .

Quadratics

As a quadratic in p , the equation is

$$(q^2 - 1)p^2 + (6q)p + (1 - 8q^2) = 0.$$

So the sum of the roots is

$$p + p' = \frac{-6q}{q^2 - 1}.$$

Quadratics

As a quadratic in p , the equation is

$$(q^2 - 1)p^2 + (6q)p + (1 - 8q^2) = 0.$$

So the sum of the roots is

$$p + p' = \frac{-6q}{q^2 - 1}.$$

As a quadratic in q , the sum of the roots is

$$q + q' = \frac{-6p}{p^2 - 8}.$$

Note that if $q = 0$, then $p = 1$. Also, if $q = 1$, then $p = \frac{7}{6}$.

An infinite chain of solutions

$$p + p' = \frac{-6q}{q^2 - 1}; \quad q + q' = \frac{-6p}{p^2 - 8}$$

If $q = 0$, then $p = 1$.

An infinite chain of solutions

$$p + p' = \frac{-6q}{q^2 - 1}; \quad q + q' = \frac{-6p}{p^2 - 8}$$

If $q = 0$, then $p = 1$. Then $q' = \frac{-6}{1 - 8} - 0 = \frac{6}{7}$.

An infinite chain of solutions

$$p + p' = \frac{-6q}{q^2 - 1}; \quad q + q' = \frac{-6p}{p^2 - 8}$$

If $q = 0$, then $p = 1$. Then $q' = \frac{-6}{1 - 8} - 0 = \frac{6}{7}$. Then

$$p' = \frac{\frac{-36}{7}}{\frac{36}{49} - 1} - 1 = \frac{239}{13} \dots$$

An infinite chain of solutions

$$p + p' = \frac{-6q}{q^2 - 1}; \quad q + q' = \frac{-6p}{p^2 - 8}$$

If $q = 0$, then $p = 1$. Then $q' = \frac{-6}{1 - 8} - 0 = \frac{6}{7}$. Then

$$p' = \frac{\frac{-36}{7}}{\frac{36}{49} - 1} - 1 = \frac{239}{13} \dots$$

Also, if $q = 1$, then $p = \frac{7}{6}$. Then $q' = \frac{-7}{\frac{49}{36} - 8} - 1 = \frac{13}{239} \dots$

An infinite chain of solutions

$$p + p' = \frac{-6q}{q^2 - 1}; \quad q + q' = \frac{-6p}{p^2 - 8}$$

If $q = 0$, then $p = 1$. Then $q' = \frac{-6}{1 - 8} - 0 = \frac{6}{7}$. Then

$$p' = \frac{\frac{-36}{7}}{\frac{36}{49} - 1} - 1 = \frac{239}{13} \dots$$

Also, if $q = 1$, then $p = \frac{7}{6}$. Then $q' = \frac{-7}{\frac{49}{36} - 8} - 1 = \frac{13}{239} \dots$

So $x = 0; \frac{6}{7}; \frac{1434}{91}; \dots$; or $x = \frac{7}{6}; \frac{91}{1434}; \dots$

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

If $x = 0$, then $\square = 1^2$.

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

If $x = 0$, then $\square = 1^2$.

If $x = \frac{6}{7}$, then $\square = \left(\frac{239}{49}\right)^2$.

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

If $x = 0$, then $\square = 1^2$.

If $x = \frac{6}{7}$, then $\square = \left(\frac{239}{49}\right)^2$.

If $x = \frac{7}{6}$, then $\square = \left(\frac{239}{36}\right)^2$.

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

If $x = 0$, then $\square = 1^2$.

If $x = \frac{6}{7}$, then $\square = \left(\frac{239}{49}\right)^2$.

If $x = \frac{7}{6}$, then $\square = \left(\frac{239}{36}\right)^2$.

If $x = \frac{1434}{91}$, then $\square = \left(\frac{2750257}{8281}\right)^2$.

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

If $x = 0$, then $\square = 1^2$.

If $x = \frac{6}{7}$, then $\square = \left(\frac{239}{49}\right)^2$.

If $x = \frac{7}{6}$, then $\square = \left(\frac{239}{36}\right)^2$.

If $x = \frac{1434}{91}$, then $\square = \left(\frac{2750257}{8281}\right)^2$.

If $x = \frac{91}{1434}$, then $\square = \left(\frac{250273387}{2056356}\right)^2$.

Check

Recall: $1 + 12x + 6x^2 + 12x^3 + x^4 = \square$.

Why is the reciprocal of a solution another solution?

If $x = 0$, then $\square = 1^2$.

If $x = \frac{6}{7}$, then $\square = \left(\frac{239}{49}\right)^2$.

If $x = \frac{7}{6}$, then $\square = \left(\frac{239}{36}\right)^2$.

If $x = \frac{1434}{91}$, then $\square = \left(\frac{2750257}{8281}\right)^2$.

If $x = \frac{91}{1434}$, then $\square = \left(\frac{250273387}{2056356}\right)^2$.

Complete?

But will this process lead to ALL the solutions to

$$a^2x^4 + 2abx^3 + cx^2 + 2bdx + d^2 = \square?$$

Complete?

But will this process lead to ALL the solutions to

$$a^2x^4 + 2abx^3 + cx^2 + 2bdx + d^2 = \square?$$

Euler goes on to exploit algebraic transformations of the original equation that allow him to find an analog of reciprocal solutions. (I.e., if the coefficients aren't symmetric, then what can be said?)

Complete?

But will this process lead to ALL the solutions to

$$a^2x^4 + 2abx^3 + cx^2 + 2bdx + d^2 = \square?$$

Euler goes on to exploit algebraic transformations of the original equation that allow him to find an analog of reciprocal solutions. (I.e., if the coefficients aren't symmetric, then what can be said?)

“...and in this way hardly any doubt can survive, but that all the satisfactory values for x will clearly be rooted out.”

Complete?

But will this process lead to ALL the solutions to

$$a^2x^4 + 2abx^3 + cx^2 + 2bdx + d^2 = 0?$$

Euler goes on to exploit algebraic transformations of the original equation that allow him to find an analog of reciprocal solutions. (I.e., if the coefficients aren't symmetric, then what can be said?)

“...and in this way hardly any doubt can survive, but that all the satisfactory values for x will clearly be rooted out.”

But is this **REALLY** all the solutions?

Thank you!

Chris Goff
University of the Pacific

cgoff@pacific.edu