



10-1-2018

NLRX1 modulates differentially NLRP3 inflammasome activation and NF- κ B signaling during *Fusobacterium nucleatum* infection

Shu Chen Hung

University of the Pacific Arthur A. Dugoni School of Dentistry

Pei Rong Huang

Chang Gung University

Cassio Luiz Coutinho Almeida-Da-Silva

University of the Pacific Arthur A. Dugoni School of Dentistry, csilva2@pacific.edu

Kalina R. Atanasova

University of Florida

Ozlem Yilmaz

Medical University of South Carolina

See next page for additional authors

Follow this and additional works at: <https://scholarlycommons.pacific.edu/dugoni-facarticles>



Part of the [Dentistry Commons](#)

Recommended Citation

Hung, S., Huang, P., Almeida-Da-Silva, C. L., Atanasova, K. R., Yilmaz, O., & Ojcius, D. M. (2018). NLRX1 modulates differentially NLRP3 inflammasome activation and NF- κ B signaling during *Fusobacterium nucleatum* infection. *Microbes and Infection*, 20(9-10), 615–625. DOI: [10.1016/j.micinf.2017.09.014](https://doi.org/10.1016/j.micinf.2017.09.014)
<https://scholarlycommons.pacific.edu/dugoni-facarticles/705>

This Article is brought to you for free and open access by the All Faculty Scholarship at Scholarly Commons. It has been accepted for inclusion in All Dugoni School of Dentistry Faculty Articles by an authorized administrator of Scholarly Commons. For more information, please contact mgibney@pacific.edu.

Authors

Shu Chen Hung, Pei Rong Huang, Cassio Luiz Coutinho Almeida-Da-Silva, Kalina R. Atanasova, Ozlem Yilmaz, and David M. Ojcius

**NLRX1 modulates differentially NLRP3 inflammasome activation
and NF- κ B signaling during *Fusobacterium nucleatum* infection**

Shu-Chen Hung ^{1,*}, Pei-Rong Huang ², Cássio Luiz Coutinho Almeida-da-Silva ^{1,3},
Kalina R. Atanasova ⁴, Ozlem Yilmaz ⁵, David M. Ojcius ^{1, §}

¹ Department of Biomedical Sciences, University of the Pacific, Arthur A. Dugoni
School of Dentistry, San Francisco, CA 94103, USA

² Center for Molecular and Clinical Immunology, Chang Gung University, Gueishan,
Taoyuan 333, Taiwan.

³ Immunobiology Program, Biophysics Institute of the Federal University of Rio
de Janeiro, RJ 21941, Brazil.

⁴ Department of Periodontology and Emerging Pathogens Institute, University of
Florida, Gainesville, FL 32610, USA

⁵ Department of Oral Health Sciences, and Department of Microbiology and
Immunology, Medical University of South Carolina, Charleston, SC, USA

* Current address: Department of Pediatrics, Stanford University, Stanford, CA
94305, USA.

§ Correspondence to: David Ojcius, University of the Pacific, Arthur Dugoni School
of Dentistry, 155 Fifth Street, San Francisco, CA 94103, USA

Email: dojcius@pacific.edu

Abstract

NOD-like receptors (NLRs) play a large role in regulation of host innate immunity, yet their role in periodontitis remains to be defined. NLRX1, a member of the NLR family that localizes to mitochondria, enhances mitochondrial ROS (mROS) generation. mROS can activate the NLRP3 inflammasome, yet the role of NLRX1 in NLRP3 inflammasome activation has not been examined. In this study, we revealed the mechanism by which NLRX1 positively regulates ATP-induced NLRP3 inflammasome activation through mROS in gingival epithelial cells (GECs). We found that depletion of NLRX1 by shRNA attenuated ATP-induced mROS generation and redistribution of the NLRP3 inflammasome adaptor protein, ASC. Furthermore, depletion of NLRX1 inhibited *Fusobacterium nucleatum* infection-activated caspase-1, suggesting that it also inhibits the NLRP3 inflammasome. Conversely, NLRX1 also acted as a negative regulator of NF- κ B signaling and IL-8 expression. Thus, NLRX1 stimulates detection of the pathogen *F. nucleatum* via the inflammasome, while dampening cytokine production. We expect that commensals should not activate the inflammasome, and NLRX1 should decrease their ability to stimulate expression of pro-inflammatory cytokines such as IL-8. Therefore, NLRX1 may act as a potential switch with regards to anti-microbial responses in healthy or diseased states in the oral cavity.

Keywords: Oral, periodontal, innate immunity, inflammation, *Fusobacterium nucleatum*

1. Introduction

F. nucleatum, a Gram-negative anaerobe, is one of the most abundant microorganisms present in the oral cavity during periodontal disease [7]. Its ability to co-aggregate with diverse microorganisms makes it a bridge that connects early commensals and late pathogens, which results in dental plaque formation [52]. In addition, *F. nucleatum* exacerbates periodontitis by inducing apoptosis of immune cells and bone loss [23, 28, 10]. However, the molecular mechanisms by which *F. nucleatum* promotes dysregulated inflammatory responses in periodontitis remain mostly unknown.

Gingival epithelial cells (GECs) represent the first protective barriers in the oral cavity that are exposed to invading microorganisms. During *F. nucleatum* infection, GECs secrete chemokines such as IL-8 to recruit neutrophils [55, 11, 26, 42]. In addition, we recently reported that GECs detect stress related to *F. nucleatum* infection by activating an inflammasome [9], which is relevant for inflammation and other physiological functions of the epithelium [36, 50], and that *F. nucleatum* infection activates the transcription factor NF- κ B and stimulates chemokine expression in GECs. However, how cells coordinate the two types of innate immune responses to *F. nucleatum* infection has not been investigated.

The innate immune system relies on germline-encoded pattern-recognition receptors (PRRs) to detect conserved pathogen-derived components that are known as pathogen-associated molecular patterns (PAMPs) or host-released factors by damaged or dying cells called damage-associated molecular patterns (DAMPs) [24, 47, 30, 48]. Based on domain homology, PRRs can be classified into 5 families. They are the Toll-like receptors (TLRs), NOD-like

1 receptors (NLRs), C-type lectin receptors (CLRs), RIG-I like receptors (RLRs), and
2 AIM2-like receptors (ALRs) [8]. TLRs and CLRs are membrane-spanning PRRs
3 detecting microorganisms in the extracellular milieu or in endosomes, whereas
4 NLRs, RLRs, and ALRs are expressed in the cytosol where they sense intracellular
5 microorganisms [8, 31].

6 NLRs are typically composed of a central conserved NOD domain, a C-
7 terminal leucine-rich repeat (LRR), and a N-terminal effector domain [45].
8 According to the N-terminal effector domain, NLRs can be further classified into
9 different groups, e.g. NLRP, NLRC, and NLRX [5]. Each NLR plays a non-redundant
10 role in recognizing specific PAMPs or DAMPs, but the responses they trigger can
11 be categorized as inflammasome-dependent or -independent. For example,
12 NLRP1, NLRP3 and NLRC4 can form inflammasomes to activate caspase-1 upon
13 sensing PAMPs and DAMPs. On the other hand, NOD1, NOD2 and NLRX1 induce
14 pro-inflammatory responses through NF- κ B or MAPK signaling during microbial
15 infection [5, 8]. NLRX1 (also known as NOD5 or NOD9) has an atypical N-terminal
16 X domain without obvious homology to other NLRs, but has a mitochondria-
17 addressing sequence that sorts NLRX1 to mitochondria [41, 56, 4].

18 Due to the localization of NLRX1 in mitochondria, its role in inflammation
19 has been linked to mitochondria, which is involved in multiple immune responses,
20 such as autophagy and inflammasome activation. Proteomic analysis has
21 identified the association between NLRX1 and mitochondrial protein TUFM (Tu
22 translation elongation factor), which interacts with autophagy-related proteins
23 Atg5-Atg12. The indirect association between NLRX1 and Atg5-Atg12 is necessary
24 to induce autophagy during viral infection in mice [33, 32]. In addition,
25 mitochondrial ROS (mROS) has been proposed to be a secondary signal for various

1 inflammatory responses. However, its generation and the mechanism by which it
2 damages pathogens remain elusive. NLRX1 can modulate mROS during infection
3 possibly through interaction with UQCRC2, a subunit in the mitochondrial
4 respiratory chain complex III [4]. Overexpression of NLRX1 enhances ROS
5 generation in response to TNF- α , poly(I:C), *Shigella* and *Chlamydia* infection,
6 whereas knockdown of NLRX1 abrogates pathogen-induced mROS generation [56,
7 2, 59]. Recently it has been shown that mROS can activate the NLRP3
8 inflammasome [60, 22, 19, 53]. Therefore, the first goal of this study is to examine
9 whether NLRX1 affects NLRP3 inflammasome activation via mROS.

10 In addition to its role in mitochondria, NLRX1 has been reported to affect
11 NF- κ B signaling [43, 57, 56]. Depletion of NLRX1 augments expression of LPS-
12 induced NF- κ B-responsive genes, such as IL-1 β and IL-6, in mice [61, 56, 3].
13 Immunoprecipitation experiments showed that NLRX1 binds constitutively to the
14 TLR downstream signaling molecule, TRAF6, under unstimulated conditions.
15 However, upon LPS stimulation, NLRX1 dissociates from TRAF6, leading to NF- κ B
16 activation [61]. In addition, TNF treatment enhances I- κ B degradation in intestinal
17 epithelial cells in conditional NLRX1 deficient mice [57]. Taken together, these
18 results show that NLRX1 may act as a negative regulator in TLR signaling.
19 Therefore, the second goal of this study is to examine the role of NLRX1 in NF- κ B
20 signaling during *Fusobacterium nucleatum* infection.

21

22

2. Materials and methods

2.1. Media, enzymes, oligonucleotides, and antibodies.

Defined Keratinocytes-serum free medium (SFM), Bovine Pituitary Extract (BPE), fetal bovine serum (FBS), TRIzol reagent, MitoSOX™ Red Mitochondrial Superoxide Indicator (M36008), JC-1 dye (T3168) and all oligonucleotides were from Thermo Fisher Scientific. ATP, puromycin, polybrene, hemin, and menadione, Triton-X, Tween-20, and paraformaldehyde were from Sigma-Aldrich. Antibodies against NLRX1 (8583), NF-κB RelA (8242) were from Cell Signaling. Anti-ASC (TMS1, ab155970 for Immunoblotting; ab64808 for microscopy) and cleaved caspase-1 p10 (ab108326) were purchased from Abcam. Anti-actin (MAB1501) and cleaved caspase-1 p20 (AB1871) were obtained from Millipore. Anti-TOPO-I (sc-5342) was purchased from Santa Cruz. Gel electrophoresis reagents were from Bio-Rad.

2.2. Cells and bacterial Culture

The human immortalized gingival epithelial cells (GECs) were obtained as previously described [22]. Cells were routinely cultured in defined keratinocyte-SFM with 5 µg/ml plasmocin and grown at 37 °C in a humidified incubator containing 5% CO₂. Cells were seeded in antibiotic-free keratinocyte-SFM medium two days before infection and during infection. An extra well was seeded for cell counting on the day of infection

F. nucleatum (ATCC 25586) was cultured anaerobically for 24 h at 37°C in brain-heart infusion broth supplemented with yeast extract (5 mg/ml), hemin (5 µg/ml), and menadione (1 µg/ml). For infection, *F. nucleatum* was cultured for 16-

24 h before harvest, followed by washing and collection in PBS. *F. nucleatum* was then added into GEC-seeded wells at an M.O.I. of 50 or 100 and incubated at 37 °C with 5% CO₂ as indicated for each assay.

2.3. Lentivirus infection and selection of shRNA-expressing Cells

GECs stably expressing shRNA against NLRX1 were generated by transducing the cells with lentiviral particles purchased from Sigma-Aldrich. The vectors encoding the 5 shRNA sequences separately were:

TRCN0000129459	NM_024618.2-1992s1c1
TRCN0000130268	NM_024618.2-3523s1c1
TRCN0000218246	NM_024618.2-1414s21c1
TRCN0000130325	NM_024618.2-3335s1c1
TRCN0000128446	NM_024618.2-3337s1c1

Transduction was carried out following the manufacturer's instructions in the presence of 8 µg/ml polybrene. Nontarget shRNA control cells were also generated using an irrelevant sequence (SHC002V, Sigma). Briefly, GEC were plated at 70% confluency 24 h prior to transduction, and the corresponding lentiviral transduction particles were added at M.O.I. of 3 overnight. Fresh media was added the next day, and stably infected cells were selected by addition of media containing 0.6 µg/ml puromycin for 1 week.

2.4. Measurement of mitochondrial membrane potential and ROS

Cells were stained with the cationic carbocyanine dye JC-1 (5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolcarbocyanine iodide) or MitoSOX mitochondrial superoxide dye as described previously [22, 21]. In brief, cells were loaded with 7.7 µM JC-1 or 5 µM MitoSOX in HBSS for 10 min at 37°C, washed with HBSS, and treated with 5 mM ATP for 20 min at 37°C followed by 3 washes with

1 HBSS. Finally, the cells were observed under wide-field fluorescence microscope
2 (Leica, Deerfield, IL). Quantitative analysis was performed using Image J software
3 from NIH.

4 5 **2.5. RNA extraction, reverse transcription, PCR and Quantitative PCR**

6 Total RNA was isolated from GECs after indicated treatments using TRIzol
7 reagent per the manufacture's protocol. Total RNA was reversed transcribed to
8 cDNA using Taqman Reverse Transcription Reagents kit (Applied Biosystem)
9 followed by PCR or qPCR as previously described [22]. The following primers
10 were used:

11 5'-GAAACCGCAGATCTCACCAT-3' and 5'-TTGTAGTGACCCGGAGGAAC-3' for
12 NLRX1; 5'-AATCTGGCAACCCTAGTCTGCTA-3' and 5'-
13 AGAAACCAAGGCACAGTGGA-3' for IL-8; 5'-TTAAAAGCAGCCCTGGTGAC-3' and
14 5'-CTCTGCTCCTCCTGTTTCGAC-3' for GAPDH; 5'-CCGCGGTAATACGTATGTCACG-3'
15 and 5'-TCCGCTTACCTCTCCAGTACTC-3' for 16S rRNA of *F. nucleatum*.

16 17 **2.6. Immunofluorescence staining**

18 For NF- κ B staining, cells infected with *F. nucleatum* were washed with
19 HBSS, fixed with 4% paraformaldehyde and permeabilized followed by 0.1%
20 Triton-X for 40 min. Following one wash with PBST (PBS containing 0.05%
21 Tween-20) and two washes with PBS, cells were blocked with 2% BSA in PBS for
22 30 min before hybridization with anti-NF- κ B p65 antibody overnight. Cells were
23 then washed with PBST once and PBS twice and incubated with goat anti-rabbit
24 IgG antibody conjugated with red fluorescent Alexa Fluor 546 dye (Invitrogen) for
25 40 min. DAPI was then used to stain nuclei for another 10 min followed by washing
26 with PBST once and PBS twice, and the fluorescence was visualized under wide-

1 field fluorescence microscope (Leica) or laser scanning confocal microscope
2 (Eclipse Ti C1, Nikon).

3 Nuclear NF- κ B p65 intensity was quantified by CellProfiler. Cells with
4 nuclear p65 mean intensity over 0.1 were considered as positive, and at least 350
5 cells were analyzed under each condition.

6 For ASC staining, the procedure is like for NF- κ B staining except the cells
7 were permeabilized for 20 min and anti-ASC and goat anti-rabbit IgG (H+L)
8 secondary antibody conjugated with Alexa Fluor 488 were used. Fluorescence
9 intensity was analyzed using ImageJ. Mean fluorescence intensity of ASC was
10 normalized to total fluorescence intensity of nuclei. At least 150 cells were
11 analyzed under each condition.

12

13 **2.7. Nuclear/cytosol fractionation and Western blots**

14 To determine NF- κ B protein levels in nuclear and cytosol, these extracts
15 were prepared using Nuclear/Cytosol Fractionation kit from BioVision. Briefly, 2
16 $\times 10^6$ uninfected or infected cells were collected and lysed with Cytosol Extraction
17 Buffer-A (CEB-A) containing DTT and protease inhibitors. Following vigorously
18 vortex and cold incubation on ice, CEB-B were added and mixed before
19 centrifugation at 16,000 g at 4°C. The supernatant (cytoplasmic extract) was
20 collected for further analysis, and the pellet was resuspended using Nuclear
21 Extraction Buffer (NEB). The mixture was vortexed for 15s every 10 min for a total
22 40 min and finally centrifuged at 16,000 g for 10 min at 4°C. The supernatant
23 (nuclear extract) was collected for future use.

24 Preparation of supernatants for caspase-1 detection and Western blot
25 were performed as previously described [22, 9].

1

2 **2.8. MTS assay**

3 MTS assay was conducted as previously described using CellTiter 9
4 AQueous One Solution Cell Proliferation Assay kit from Promega [21].

5

6 **2.9. Statistical analysis**

7 Immunoblot and fluorescence microscopy data were quantitatively analyzed
8 using ImageJ software or CellProfiler software. Statistics were carried out using two-
9 way ANOVA with Bonferroni's *post hoc* test. For experiments examining knockdown
10 efficiency, one-way ANOVA was used. A p value less than 0.05 was considered
11 significant.

12

13

3. Results

3.1. Attenuated ATP-induced mROS in NLRX1-deficient GECs

We have previously reported that a DAMP, extracellular ATP, induces mROS generation to activate the NLRP3 inflammasome in GECs [22]. However, the mechanism by which mROS were generated remained to be investigated. NLRX1, the only NLR localized in mitochondria, has been shown to mediate mROS production during microbial infection [56, 2, 59]. Therefore, we depleted GECs of NLRX1 using shRNA specific against NLRX1 or tGFP as a control. qRT-PCR and Western blot confirmed that NLRX1 was efficiently depleted in GECs transduced with shRNA sequence 325 and 446, compared with the tGFP control (Fig. 1, A and B).

We next measured ATP-induced mROS generation in cells stained with MitoSOX, which specifically detects mitochondrial superoxide. Fluorescence microscopy showed that ATP stimulated a large increase in mROS production in wild-type GECs and GECs transduced with shRNA control, but lower levels of mROS were observed in the NLRX1-depleted cells. Quantification analysis of the fluorescence images showed that ATP-promoted ROS production decreased ~25% in NLRX1-depleted cells compared with wild type cells (Fig. 1 C and D). These results suggest that NLRX1 enhances ATP-triggered mROS generation.

3.2. Depletion of NLRX1 reduced ATP-induced mitochondrial damage

Recent studies revealed a critical role of mitochondria in NLRP3 inflammasome activation [53, 64]. Thus, NLRP3 agonists such as ATP disrupt the mitochondrial membrane potential, which results in mROS generation and

1 subsequent release of oxidized mitochondrial DNA to the cytosol, where it
2 activates the NLRP3 inflammasome [53]. We therefore examined the effect of
3 NLRX1 on mitochondrial membrane potential ($\Delta\psi_m$) in GECs with the
4 mitochondrial membrane potential probe, JC-1. The dye JC-1 shifts profoundly
5 from red to green in both WT and shCtrl GECs upon ATP stimulation, suggesting
6 that ATP triggered a loss in $\Delta\psi_m$. However, depletion of NLRX1 reduced the ATP-
7 elicited $\Delta\psi_m$ decrease (Fig. 2A). Quantitative analysis of the fluorescence
8 micrographs confirmed that depletion of NLRX1 abrogates the ATP-induced $\Delta\psi_m$
9 decrease (Fig. 2B).

10 ATP-induced mROS also induces the cytosolic translocation of
11 mitochondrial DNA (mtDNA), which is recognized by the NLRP3 inflammasome
12 [40, 53]. We therefore examined whether NLRX1 had an effect on ATP-triggered
13 mtDNA release to cytosol in GECs. Cytosol were separated and collected from ATP-
14 treated or untreated cells to measure mitochondrial DNA encoding *Cyt b* by PCR.
15 Incubation with ATP significantly increased the level of cytosolic mtDNA in GECs,
16 whereas NLRX1 depletion abrogated the enhancement (Fig. 2C and 2C). Genomic
17 DNA encoding *Gapdh* was not detected in the cytosolic samples, excluding the
18 possibility of nonspecific contamination from organelles. Taken together, these
19 results suggest that NLRX1 protects against ATP-caused mitochondrial damage.

20

21 **3.3. Diminished NLRP3 activation in NLRX- depleted GECs**

22 Mitochondrial dysfunction-caused cytosolic translocation of mitochondrial
23 DNA is sensed by NLRP3, which triggers NLRP3 inflammasome assembly [53].
24 Assembly of the NLRP3 inflammasome requires recruitment of the adaptor
25 protein ASC (apoptosis-associated speck-like protein containing CARD) at

1 perinuclear mitochondria-associated membranes (MAMs) of the endoplasmic
2 reticulum (ER) [64, 38, 35]. To determine whether NLRX1 is involved in the
3 redistribution of ASC, the cells were treated with ATP for 10 min followed by
4 immunofluorescence assays. Under untreated conditions, ASC was barely
5 detectable due to its general distribution throughout the cytosol. However, in
6 ATP-stimulated GECs with or without shRNA against tGFP, ASC became
7 predominantly perinuclear, and NLRX1-depletion blocked the ATP-induced ASC
8 redistribution (Fig. 3A, 3B, and Fig. S1). In addition, the protein abundance of ASC
9 was not altered by the treatment (Fig. 3C). We also examined whether NLRX1
10 contributes to *F. nucleatum* infection-triggered ASC redistribution. As Fig. S2
11 shows, *F. nucleatum* infection leads to ASC to become concentrated to in the
12 perinuclear region, consistent with our previous observation that NLRP3
13 inflammasome is activated by infection [9]. NLRX1 depletion also suppressed the
14 perinuclear relocalization of ASC by infection (Fig. S2). Taken together, these data
15 suggest that NLRX1 abolishes ATP- and *F. nucleatum*- triggered ASC redistribution
16 rather than affecting ASC expression.

17 NLRP3 inflammasome assembly with ASC leads to autoprocessing and
18 activation of caspase-1, which in turn cleaves pro-IL-1 β to IL-1 β [1]. To determine
19 whether the compromised redistribution of ASC affected NLRP3 inflammasome
20 activation, we measured the protein level of activated caspase-1 by Western blots.
21 As expected, ATP stimulation significantly enhanced caspase-1 activation in wt
22 and shCtrl GECs. The level of caspase-1 was also enhanced in NLRX1-depleted cells
23 after ATP stimulation, but to a lesser extent than in wild-type cells. Caspase-1
24 activation was impaired more severely in cells transduced with shNLRX1
25 sequence 446, which may be due to the better knockdown efficiency of NLRX1

1 with this sequence (Fig. 3D). Taken together, these data suggest that NLRX1
2 regulates positively NLRP3 inflammasome activation.

3 4 **3.4. NLRX1 enhances caspase-1 activation during *F. nucleatum* infection**

5 Previous studies have shown that *F. nucleatum* infection activates the
6 NLRP3 inflammasome [58, 9]. Since NLRX1 modulates ATP-induced NLRP3
7 inflammasome activation, we hypothesized that NLRX1 is also involved in *F.*
8 *nucleatum* mediated NLRP3 inflammasome activation. We therefore examined
9 caspase-1 activation during infection of GECs by *F. nucleatum*, and observed that
10 infection-induced caspase-1 activation decreased in NLRX1-deficient cells,
11 compared with controls. Cell viability was not affected by the depletion of NLRX1
12 and was not significantly affected by infection (Fig. 4 and Fig. S3) [9] [25, 15].
13 Taken together, these results suggest that NLRX1 positively modulates *F.*
14 *nucleatum* infection-elicited NLRP3 inflammasome activation.

15 16 **3.5. Enhanced IL-8 gene expression in NLRX1-deficient cells during *F.*** 17 ***nucleatum* infection**

18 IL-8 is expressed via NF- κ B signaling to recruit neutrophils in response to
19 *F. nucleatum* infection [11, 63]. However, the mechanism whereby *F. nucleatum*
20 increases IL-8 expression remains elusive. NLRX1 has been reported to negatively
21 regulate NF- κ B signaling by targeting the IKK complex [61]. Therefore, we
22 examined whether NLRX1 participates in *F. nucleatum*-induced IL-8 expression.
23 Cells were infected with or without *F. nucleatum* at a multiplicity of infection
24 (M.O.I.) of 50 and 100, and IL-8 expression was then measured by qPCR. As shown
25 in Fig. 5A, IL-8 gene expression increased as a function of *F. nucleatum* dose in

1 shCtrl and wildtype cells, but infection in NLRX1-deficient cells led to significantly
2 higher IL-8 expression. 16S rRNA gene was examined by RT-PCR to confirm
3 infection (Fig. 5B). These data suggest that NLRX1 negatively regulates *F.*
4 *nucleatum* infection-induced IL-8 expression.

6 **3.6. NLRX1 modulates IL-8 expression during *F. nucleatum* infection via the** 7 **transcription factor NF- κ B**

8 NLRX1 has been previously reported to negatively regulate NF- κ B
9 signaling [61]. To determine whether NLRX1 is involved in *F. nucleatum* infection-
10 induced IL-8 expression, the nuclear translocation of RelA (p65) subunit of NF- κ B
11 was examined in NLRX1 knockdown GECs by confocal microscopy. As expected,
12 infection by *F. nucleatum* resulted in nuclear translocation of RelA in 30% of
13 wildtype and shCtrl cells, but RelA translocation increased to 60% in NLRX1-
14 deficient cells (Fig. 6A, 6B and Fig. S4). In addition, the cytosolic and nuclear
15 extracts from infected cells were analyzed for RelA by immunoblotting. NLRX1-
16 deficient cells showed increased RelA levels in the nucleus, but reduced RelA in
17 the cytosol during infection, compared to wildtype and shCtrl cells (Fig. 6C). These
18 results suggest that NLRX1 negatively regulates IL-8 expression through its action
19 on NF- κ B.

20
21
22

4. Discussion

The role of NLRX1 in innate immunity is still under investigation. Initial studies showed that NLRX1 negatively regulates RIG-I-MAVS-dependent IFN- β expression to prevent excessive inflammation during viral infection [62, 3, 43]. In addition, SIV infection leads to upregulation of NLRX1 and downregulation of interferon-stimulated genes (ISGs) at early stages of infection to promote viral dissemination in rhesus monkeys [6]. More recent studies demonstrated that NLRX1 attenuates STING-dependent ISGs upon HIV-1 or DNA virus infection in human macrophages and dendritic cells [14]. However, the role of NLRX1 in immune responses to bacterial infection remains less clear.

F. nucleatum induces NF- κ B-mediated IL-8 expression in GECs, which results in neutrophil recruitment to eradicate infection [20, 63]. However, the pattern recognition receptors (PRRs) responsible for sensing the bacteria remain elusive. IL-8 is expressed following stimulation of extracellular Toll-like receptors (TLRs) during most bacterial infections. Nevertheless, it has been reported that some *F. nucleatum* strains upregulate IL-8 through TLR-independent mechanisms [63, 34]. Transcriptome analysis and a study using TLR-deficient HEK293T cells have shown that mitogen-activated protein kinase (MAPK) signaling also contribute to IL-8 upregulation upon *F. nucleatum* infection [17, 46]. The ability of *F. nucleatum* to invade cells suggests that intracellular or endosomal PRRs may also contribute to IL-8 production [46, 16]. Consistent with this possibility, it has been shown that endosomal TLR9 up-regulates IL-8 in response to *F. nucleatum* DNA in GECs [29]. Furthermore, the crystal structure of NLRX1 suggests that its leucine-rich repeat (LRR) domain could directly sense PAMPs, such as nucleic acids [18]. Here, we demonstrated that depletion of NLRX1 enhanced RelA nuclear translocation, thus

1 augmenting IL-8 expression. Previous research showing that NLRX1 negatively regulates
2 the NF- κ B pathway through suppression of TRAF6 is consistent with our results [3, 43,
3 61]. However, IL-8 can also be regulated by the MAPK pathway, which is also downstream
4 from TRAF6 [37]. Therefore, further study is needed to validate whether NLRX1
5 modulates *F. nucleatum*-triggered IL-8 expression via the NF- κ B pathway alone.

6 We recently reported that *F. nucleatum* infection activates the NLRP3
7 inflammasome and facilitates the release of DAMPs such as HMGB1 and ASC to amplify
8 inflammation in GECs [9]. Here, we have demonstrated that NLRX1 behaves as a positive
9 regulator in *F. nucleatum* infection-elicited NLRP3 inflammasome activation. Consistent
10 with our finding, it has been proposed that NLRX1 modulates NLRP3 inflammasome
11 activation during *Chlamydia* infection [13, 2].

12 DAMPs released from infected or stressed cells, such as extracellular ATP, could
13 activate the NLRP3 inflammasome through P2X₄/P2X₇-gated Pannexin-1 channels, which
14 promote mROS generation [22]. Previous reports identified a role for NLRX1 in
15 enhancing ROS production during *Shigella* infection [56, 61]. Thus, internalized *F.*
16 *nucleatum* could destabilize host-cell mitochondria, which results in mROS production,
17 which in turn is regulated by NLRX1 [54, 39, 51, 12, 27]. Interestingly, a recent study
18 revealed up-regulation of NLRP3 expression but down-regulation of NLRX1 in
19 *Helicobacter pylori*-infected bone-marrow derived macrophages [44, 49].

20 In this study, we uncovered a surprising role for NLRX1 as a molecular switch
21 between NLRP3 inflammasome activation and NF- κ B signaling in GECs during *F.*
22 *nucleatum* infection. Depletion of NLRX1 decreases NLRP3 inflammasome activation due
23 to ATP stimulation and *F. nucleatum* infection. Therefore, we hypothesize that NLRX1
24 promotes dysregulated inflammatory responses during *F. nucleatum* infection via the
25 NLRP3 inflammasome in periodontitis. Conversely, commensals or *F. nucleatum* in

1 healthy individuals may still induce NF- κ B-mediated IL-8 expression, but it is repressed
2 by NLRX1. Thus, we propose that NLRX1 should enhance the innate immune response
3 during infection by pathogens but behave as a break to prevent excessive inflammation
4 under normal circumstances (Fig. 7).

5

6 **Conflict of interest**

7 The authors declare no conflicts of interest.

8

9 **Acknowledgements**

10 This work was supported by funds from the University of the Pacific, Fundação de
11 Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) in Brazil, and the National
12 Institutes of Health grant R01 DE016593.

13

5. Reference

- [1] Abderrazak A, Syrovets T, Couchie D, El Hadri K, Friguet B, Simmet T, et al. NLRP3 inflammasome: from a danger signal sensor to a regulatory node of oxidative stress and inflammatory diseases. *Redox Biol* 2015;4:296-307.
- [2] Abdul-Sater AA, Said-Sadier N, Lam VM, Singh B, Pettengill MA, Soares F, et al. Enhancement of reactive oxygen species production and chlamydial infection by the mitochondrial Nod-like family member NLRX1. *J Biol Chem* 2010;285:41637-45.
- [3] Allen IC, Moore CB, Schneider M, Lei Y, Davis BK, Scull MA, et al. NLRX1 protein attenuates inflammatory responses to infection by interfering with the RIG-I-MAVS and TRAF6-NF-kappaB signaling pathways. *Immunity* 2011;34:854-65.
- [4] Arnoult D, Soares F, Tattoli I, Castanier C, Philpott DJ, Girardin SE. An N-terminal addressing sequence targets NLRX1 to the mitochondrial matrix. *J Cell Sci* 2009;122:3161-8.
- [5] Barbe F, Douglas T, Saleh M. Advances in Nod-like receptors (NLR) biology. *Cytokine Growth Factor Rev* 2014;25:681-97.
- [6] Barouch DH, Ghneim K, Bosche WJ, Li Y, Berkemeier B, Hull M, et al. Rapid Inflammasome Activation following Mucosal SIV Infection of Rhesus Monkeys. *Cell* 2016;165:656-67.
- [7] Bolstad AI, Jensen HB, Bakken V. Taxonomy, biology, and periodontal aspects of *Fusobacterium nucleatum*. *Clin Microbiol Rev* 1996;9:55-71.
- [8] Brubaker SW, Bonham KS, Zanoni I, Kagan JC. Innate immune pattern recognition: a cell biological perspective. *Annu Rev Immunol* 2015;33:257-90.
- [9] Bui FQ, Johnson L, Roberts J, Hung SC, Lee J, Atanasova KR, et al. *Fusobacterium nucleatum* infection of gingival epithelial cells leads to NLRP3 inflammasome-dependent secretion of IL-1beta and the danger signals ASC and HMGB1. *Cell Microbiol* 2016;18:970-81.
- [10] Chaushu S, Wilensky A, Gur C, Shapira L, Elboim M, Halftek G, et al. Direct recognition of *Fusobacterium nucleatum* by the NK cell natural cytotoxicity receptor NKp46 aggravates periodontal disease. *PLoS Pathog* 2012;8:e1002601.
- [11] Darveau RP, Belton CM, Reife RA, Lamont RJ. Local chemokine paralysis, a novel pathogenic mechanism for *Porphyromonas gingivalis*. *Infect Immun* 1998;66:1660-5.
- [12] Duncan JA, Gao X, Huang MT, O'Connor BP, Thomas CE, Willingham SB, et al. *Neisseria gonorrhoeae* activates the proteinase cathepsin B to mediate the signaling activities of the NLRP3 and ASC-containing inflammasome. *J Immunol* 2009;182:6460-9.

- 1 [13] Elwell C, Mirrashidi K, Engel J. *Chlamydia* cell biology and pathogenesis. Nat
2 Rev Microbiol 2016;14:385-400.
- 3 [14] Guo H, Konig R, Deng M, Riess M, Mo J, Zhang L, et al. NLRX1 Sequesters
4 STING to Negatively Regulate the Interferon Response, Thereby Facilitating the
5 Replication of HIV-1 and DNA Viruses. Cell Host Microbe 2016;19:515-28.
- 6 [15] Gursoy UK, Kononen E, Uitto VJ. Intracellular replication of fusobacteria
7 requires new actin filament formation of epithelial cells. APMIS 2008;116:1063-
8 70.
- 9 [16] Han YW, Shi W, Huang GT, Kinder Haake S, Park NH, Kuramitsu H, et al.
10 Interactions between periodontal bacteria and human oral epithelial cells:
11 *Fusobacterium nucleatum* adheres to and invades epithelial cells. Infect Immun
12 2000;68:3140-6.
- 13 [17] Hasegawa Y, Mans JJ, Mao S, Lopez MC, Baker HV, Handfield M, et al. Gingival
14 epithelial cell transcriptional responses to commensal and opportunistic oral
15 microbial species. Infect Immun 2007;75:2540-7.
- 16 [18] Hong M, Yoon SI, Wilson IA. Structure and functional characterization of the
17 RNA-binding element of the NLRX1 innate immune modulator. Immunity
18 2012;36:337-47.
- 19 [19] Horng T. Calcium signaling and mitochondrial destabilization in the
20 triggering of the NLRP3 inflammasome. Trends Immunol 2014;35:253-61.
- 21 [20] Huang GT, Zhang HB, Dang HN, Haake SK. Differential regulation of cytokine
22 genes in gingival epithelial cells challenged by *Fusobacterium nucleatum* and
23 *Porphyromonas gingivalis*. Microb Pathog 2004;37:303-12.
- 24 [21] Huang PR, Hung SC, Pao CC, Wang TC. N-(1-pyrenyl) maleimide induces bak
25 oligomerization and mitochondrial dysfunction in Jurkat Cells. Biomed Res Int
26 2015;2015:798489.
- 27 [22] Hung SC, Choi CH, Said-Sadier N, Johnson L, Atanasova KR, Sellami H, et al.
28 P2X4 assembles with P2X7 and pannexin-1 in gingival epithelial cells and
29 modulates ATP-induced reactive oxygen species production and inflammasome
30 activation. PLoS One 2013;8:e70210.
- 31 [23] Huynh T, Kapur RV, Kaplan CW, Cacalano N, Kinder Haake S, Shi W, et al.
32 The role of aggregation in *Fusobacterium nucleatum*- induced immune cell death.
33 J Endod 2011;37:1531-5.
- 34 [24] Janeway CA, Jr., Medzhitov R. Innate immune recognition. Annu Rev
35 Immunol 2002;20:197-216.
- 36 [25] Ji S, Shin JE, Kim YC, Choi Y. Intracellular degradation of *Fusobacterium*
37 *nucleatum* in human gingival epithelial cells. Mol Cells 2010;30:519-26.

- 1 [26] Ji S, Shin JE, Kim YS, Oh JE, Min BM, Choi Y. Toll-like receptor 2 and NALP2
2 mediate induction of human beta-defensins by *Fusobacterium nucleatum* in
3 gingival epithelial cells. Infect Immun 2009;77:1044-52.
- 4 [27] Jo EK, Kim JK, Shin DM, Sasakawa C. Molecular mechanisms regulating
5 NLRP3 inflammasome activation. Cell Mol Immunol 2016;13:148-59.
- 6 [28] Kaplan CW, Lux R, Huynh T, Jewett A, Shi W, Haake SK. *Fusobacterium*
7 *nucleatum* apoptosis-inducing outer membrane protein. J Dent Res 2005;84:700-
8 4.
- 9 [29] Kim Y, Jo AR, Jang da H, Cho YJ, Chun J, Min BM, et al. Toll-like receptor 9
10 mediates oral bacteria-induced IL-8 expression in gingival epithelial cells.
11 Immunol Cell Biol 2012;90:655-63.
- 12 [30] Krysko DV, Agostinis P, Krysko O, Garg AD, Bachert C, Lambrecht BN, et al.
13 Emerging role of damage-associated molecular patterns derived from
14 mitochondria in inflammation. Trends Immunol 2011;32:157-64.
- 15 [31] Kumar S, Ingle H, Prasad DV, Kumar H. Recognition of bacterial infection by
16 innate immune sensors. Crit Rev Microbiol 2013;39:229-46.
- 17 [32] Lei Y, Wen H, Ting JP. The NLR protein, NLRX1, and its partner, TUFM,
18 reduce type I interferon, and enhance autophagy. Autophagy 2013;9:432-3.
- 19 [33] Lei Y, Wen H, Yu Y, Taxman DJ, Zhang L, Widman DG, et al. The
20 mitochondrial proteins NLRX1 and TUFM form a complex that regulates type I
21 interferon and autophagy. Immunity 2012;36:933-46.
- 22 [34] Liu H, Redline RW, Han YW. *Fusobacterium nucleatum* induces fetal death in
23 mice via stimulation of TLR4-mediated placental inflammatory response. J
24 Immunol 2007;179:2501-8.
- 25 [35] Man SM, Hopkins LJ, Nugent E, Cox S, Gluck IM, Tourlomousis P, et al.
26 Inflammasome activation causes dual recruitment of NLRC4 and NLRP3 to the
27 same macromolecular complex. Proc Natl Acad Sci U S A 2014;111:7403-8.
- 28 [36] Martel J, Lai HC, Ko YF, Young JD, Ojcius DM. Alternative functions for the
29 multifarious inflammasome. Biomed J 2016;39:183-7.
- 30 [37] McClure R, Massari P. TLR-Dependent Human Mucosal Epithelial Cell
31 Responses to Microbial Pathogens. Front Immunol 2014;5:386.
- 32 [38] Misawa T, Takahama M, Kozaki T, Lee H, Zou J, Saitoh T, et al. Microtubule-
33 driven spatial arrangement of mitochondria promotes activation of the NLRP3
34 inflammasome. Nat Immunol 2013;14:454-60.
- 35 [39] Murakami T, Ockinger J, Yu J, Byles V, McColl A, Hofer AM, et al. Critical role
36 for calcium mobilization in activation of the NLRP3 inflammasome. Proc Natl
37 Acad Sci U S A 2012;109:11282-7.

- 1 [40] Nakahira K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, Lam HC, et al.
2 Autophagy proteins regulate innate immune responses by inhibiting the release
3 of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol*
4 2011;12:222-30.
- 5 [41] O'Neill LA. Innate immunity: squelching anti-viral signalling with NLRX1.
6 *Curr Biol* 2008;18:R302-4.
- 7 [42] Park SR, Kim DJ, Han SH, Kang MJ, Lee JY, Jeong YJ, et al. Diverse Toll-like
8 receptors mediate cytokine production by *Fusobacterium nucleatum* and
9 *Aggregatibacter actinomycetemcomitans* in macrophages. *Infect Immun*
10 2014;82:1914-20.
- 11 [43] Parvatiyar K, Cheng G. NOD so fast: NLRX1 puts the brake on inflammation.
12 *Immunity* 2011;34:821-2.
- 13 [44] Philipson CW, Bassaganya-Riera J, Viladomiu M, Kronsteiner B, Abedi V,
14 Hoops S, et al. Modeling the Regulatory Mechanisms by Which NLRX1 Modulates
15 Innate Immune Responses to *Helicobacter pylori* Infection. *PLoS One*
16 2015;10:e0137839.
- 17 [45] Proell M, Riedl SJ, Fritz JH, Rojas AM, Schwarzenbacher R. The Nod-like
18 receptor (NLR) family: a tale of similarities and differences. *PLoS One*
19 2008;3:e2119.
- 20 [46] Quah SY, Bergenholtz G, Tan KS. *Fusobacterium nucleatum* induces cytokine
21 production through Toll-like-receptor-independent mechanism. *Int Endod J*
22 2014;47:550-9.
- 23 [47] Rubartelli A, Lotze MT. Inside, outside, upside down: damage-associated
24 molecular-pattern molecules (DAMPs) and redox. *Trends Immunol* 2007;28:429-
25 36.
- 26 [48] Said-Sadier N, Ojcius DM. Alarmins, inflammasomes and immunity. *Biomed J*
27 2012;35:437-49.
- 28 [49] Salama NR, Hartung ML, Muller A. Life in the human stomach: persistence
29 strategies of the bacterial pathogen *Helicobacter pylori*. *Nat Rev Microbiol*
30 2013;11:385-99.
- 31 [50] Santana PT, Martel J, Lai HC, Perfettini JL, Kanellopoulos JM, Young JD, et al.
32 Is the inflammasome relevant for epithelial cell function? *Microbes Infect*
33 2016;18:93-101.
- 34 [51] Shao BZ, Xu ZQ, Han BZ, Su DF, Liu C. NLRP3 inflammasome and its
35 inhibitors: a review. *Front Pharmacol* 2015;6:262.
- 36 [52] Sharma A, Inagaki S, Sigurdson W, Kuramitsu HK. Synergy between
37 *Tannerella forsythia* and *Fusobacterium nucleatum* in biofilm formation. *Oral*
38 *Microbiol Immunol* 2005;20:39-42.

- 1 [53] Shimada K, Crother TR, Karlin J, Dagvadorj J, Chiba N, Chen S, et al. Oxidized
2 mitochondrial DNA activates the NLRP3 inflammasome during apoptosis.
3 Immunity 2012;36:401-14.
- 4 [54] Shin JE, Baek KJ, Choi YS, Choi Y. A periodontal pathogen *Treponema*
5 *denticola* hijacks the *Fusobacterium nucleatum*-driven host response. Immunol
6 Cell Biol 2013;91:503-10.
- 7 [55] Signat B, Roques C, Poulet P, Duffaut D. *Fusobacterium nucleatum* in
8 periodontal health and disease. Curr Issues Mol Biol 2011;13:25-36.
- 9 [56] Tattoli I, Carneiro LA, Jehanno M, Magalhaes JG, Shu Y, Philpott DJ, et al.
10 NLRX1 is a mitochondrial NOD-like receptor that amplifies NF-kappaB and JNK
11 pathways by inducing reactive oxygen species production. EMBO Rep
12 2008;9:293-300.
- 13 [57] Tattoli I, Killackey SA, Foerster EG, Molinaro R, Maisonneuve C, Rahman MA,
14 et al. NLRX1 Acts as an Epithelial-Intrinsic Tumor Suppressor through the
15 Modulation of TNF-Mediated Proliferation. Cell Rep 2016;14:2576-86.
- 16 [58] Taxman DJ, Swanson KV, Broglie PM, Wen H, Holley-Guthrie E, Huang MT, et
17 al. *Porphyromonas gingivalis* mediates inflammasome repression in
18 polymicrobial cultures through a novel mechanism involving reduced
19 endocytosis. J Biol Chem 2012;287:32791-9.
- 20 [59] Unger BL, Ganesan S, Comstock AT, Faris AN, Hershenson MB, Sajjan US.
21 Nod-like receptor X-1 is required for rhinovirus-induced barrier dysfunction in
22 airway epithelial cells. J Virol 2014;88:3705-18.
- 23 [60] Weinberg SE, Sena LA, Chandel NS. Mitochondria in the regulation of innate
24 and adaptive immunity. Immunity 2015;42:406-17.
- 25 [61] Xia X, Cui J, Wang HY, Zhu L, Matsueda S, Wang Q, et al. NLRX1 negatively
26 regulates TLR-induced NF-kappaB signaling by targeting TRAF6 and IKK.
27 Immunity 2011;34:843-53.
- 28 [62] Xiao TS, Ting JP. NLRX1 has a tail to tell. Immunity 2012;36:311-2.
- 29 [63] Zhang G, Chen R, Rudney JD. Streptococcus cristatus modulates the
30 *Fusobacterium nucleatum*-induced epithelial interleukin-8 response through the
31 nuclear factor-kappa B pathway. J Periodontal Res 2011;46:558-67.
- 32 [64] Zhou R, Yazdi AS, Menu P, Tschopp J. A role for mitochondria in NLRP3
33 inflammasome activation. Nature 2011;469:221-5.
- 34
- 35

Figure Legends

Fig. 1. Depletion of NLRX1 reduced ATP-induced mROS in GECs.

GECs were transduced with lentivirus carrying specific shRNA targeting NLRX1 (A) NLRX1 mRNA levels were measured by qRT-PCR in GECs expressing different shRNA. (B) Whole cell lysates were collected to measure protein expression levels of NLRX1. Actin was used as an internal control. (C and D) mROS generation was detected by fluorescence microscopy using MitoSOX after 5 mM ATP stimulation for 20 min. Quantification of mROS production were analyzed by ImageJ as in (C). BF represents bright field. Mean fluorescence intensity was obtained by dividing total fluorescence intensity against total number of cells in each image. Relative mean fluorescence is determined by normalizing with control group in each cell lines. Scale bars represents 100 μ m. Statistical significance determined using one-way ANOVA for (A) and two-way ANOVA for (C). *P-value < 0.05; **P-value < 0.01; ***P-value < 0.001. Data were from a representative of at least three independent experiments performed in duplicates or triplicates that gave similar results.

Fig. 2. Suppression of ATP-elicited mitochondrial dysfunction in NLRX1-depleted GECs.

(A) GECs depleted of NLRX1 were stained with 5 μ g/ml JC-1 followed by 5 mM ATP treatment for 20 min. The treated cells were subject to fluorescence microscopy for the analysis of the loss of mitochondrial membrane potential ($\Delta\Psi_m$). Red indicates JC-1 aggregates in healthy cells with high membrane potential. The red aggregates shifts to green monomer in cytosol when mitochondrial potential is depolarized. Scale bars: 100 μ m. (B) Quantification of

1 mitochondrial membrane depolarization was obtained by dividing total
2 fluorescence intensity of red J-aggregate over green monomer followed by
3 normalization to the PBS control group. Data shown are representative of two or
4 more independent experiments performed in duplicates (means \pm SD). Statistical
5 significance was determined using two-way ANOVA. ***P-value < 0.001. (C)
6 Cytosolic fractions were isolated from GECs treated with or without 5 mM ATP for
7 1 h followed by Phenol-Chloroform-Isoamyl Alcohol precipitation. The cytosolic
8 DNA were subject to PCR to detect mitochondrial gene CytB and genomic GAPDH.
9 Data shown are representative of two or more independent experiments (means \pm
10 SD).

11

12 **Fig. 3. NLRX1-depletion led to lower ATP-mediated NLRP3 inflammasome**
13 **activation.**

14 (A) GECs were treated or untreated with 5 mM ATP for 10 min before fixation. The
15 cells were then immunostained for ASC (green) antibody and subjected to
16 confocal microscopy. Nuclei were stained by DAPI (blue). Scale bar, 20 μ m. (B)
17 Fluorescence microscopy images were quantified by dividing mean ASC
18 fluorescence intensity over total DAPI fluorescence intensity of each cell.
19 Statistical significance was determined using two-way ANOVA. **P-value < 0.01;
20 ***P-value = 0.001. (C) Western blot analysis of ASC protein levels of whole cell
21 lysates from GECs treated as in (A). Actin was chosen as loading control. (D)
22 Caspase-1 protein levels were examined by Western blot after GECs were left
23 untreated or treated with 5 mM ATP for 1 h. Actins was used as internal control.
24 Data shown are representative of two or more independent experiments (means \pm
25 SD).

Fig. 4. NLRX1 depletion decreased *F. nucleatum*-triggered caspase-1 activation.

GECs were infected with or without *F. nucleatum* (Fn) at an MOI of 100 for 24 h. Supernatants were collected and subjected to TCA precipitation. Cells were lysed in RIPA buffer. Western blot was assayed to determine the protein levels of caspase-1 in supernatants (extracellular) and whole cell lysates (intracellular). Actin served as loading control. Data are representative of three independent experiments.

Fig. 5. NLRX1 depletion enhanced IL-8 expression upon *F. nucleatum* infection.

(A) GECs were left uninfected or infected with *F. nucleatum* at an M.O.I. of 50 for 24 h. RNA extracted from the cells were reverse transcribed to cDNA followed by qPCR to detect the IL-8 transcripts. Statistical significance was determined using two-way ANOVA. ***P-value < 0.001. (B) PCR were used to detect 16s rRNA of *F. nucleatum* to confirm the infection in (A). GAPDH was used as an internal control. Data are representative of three independent experiments performed in triplicates (means \pm SD).

Fig. 6. NLRX1 regulated *F. nucleatum*-elicited NF- κ B nuclear translocation.



(A) Confocal microscopy of GECs uninfected or infected with *F. nucleatum* at an M.O.I. of 50 for 1h followed by fixation. Cells were then immunostained with anti-NF- κ B p65 (red) and stained nuclei with DAPI (blue). Scale bar, 20 μ m. (B) Quantitative analysis of fluorescence microscopy images obtained under

1 condition as in (A). Fluorescence intensity of p65 in nuclei and total cells was
2 measured by CellProfiler. Statistical significance was determined using two-way
3 ANOVA. *** $P < 0.001$. (C) Nuclear and cytosol fraction of cells were isolated
4 followed by immunoblotting using anti-NF- κ B p65 antibody. TOPO I and actin
5 served as loading control for nuclear and cytosol fractions, separately. All data are
6 representative of at least three independent experiments performed in duplicates.


7

8 **Fig. 7.** Model of *F. nucleatum* infection-triggered inflammatory responses in
9 gingival epithelial cells during healthy (dotted line) or diseased status (solid line).
10 Pathogens and danger signals (DS) such as ATP activate the NLRP3 inflammasome,
11 which is enhanced by NLRX1. Bacteria under non-pathogenic conditions can also
12 stimulate NF- κ B activation, which is dampened by NLRX1.


13




(A)




(B)




(C)




(A)




(B)




(C)




(D)






(A)






(B)




(A)

Uninfected

Fn

(B)



(C)

